



Original Article

Towards spatial management of fisheries in the Gulf: benthic diversity, habitat and fish distributions from Qatari waters

Mark E. M. Walton^{1,*}, Jamie Hayes¹, Mohsin Al-Ansi², Mohamed Abdallah³, Ibrahim Al Maslamani⁴, Mohammed Al-Mohannadi³, Ismail Al-Shaikh^{4,5}, Tim D'Urban Jackson¹, Claire Szostek¹, Jack Egerton⁶, Michel J. Kaiser⁶, and Lewis Le Vay¹

¹Centre for Applied Marine Sciences, Bangor University, Menai Bridge, Wales, UK

²Department of Biology, Qatar University, Doha, State of Qatar

³Department of Fisheries, Ministry of Municipality and Environment, Doha, State of Qatar

⁴Environmental Science Center, Qatar University, Doha, State of Qatar

⁵ExxonMobil Research Qatar, Qatar Science and Technology Park – Tech 2, PO Box 22500, Doha, State of Qatar

⁶School of Ocean Sciences, Bangor University, Menai Bridge, Wales, UK

*Corresponding author: e-mail: m.walton@bangor.ac.uk

Walton, M. E. M., Hayes, J., Al-Ansi, M., Abdallah, M., Al Maslamani, I., Al-Mohannadi, M., Al-Shaikh, I., D'Urban Jackson, T., Szostek, C., Egerton, J., Kaiser, M. J., and Le Vay, L. Towards spatial management of fisheries in the Gulf: benthic diversity, habitat and fish distributions from Qatari waters. – ICES Journal of Marine Science, 75: 178–189.

Received 21 March 2017; revised 12 May 2017; accepted 21 May 2017; advance access publication 27 July 2017.

As with many other regions in the world, more complete information on the distribution of marine habitats in the Gulf is required to inform environmental policy, and spatial management of fisheries resources will require better understanding of the relationships between habitat and fish communities. Towed cameras and sediment grabs were used to investigate benthic habitats and associated epifauna, infauna and fish communities in the central Gulf, offshore from the east coast of Qatar, in water depths of between 12 and 52 m. Six different habitats were identified: (i) soft sediment habitats of mud and (ii) sand, and structured habitats of (iii) macro-algal reef, (iv) coral reef, (v) mixed reef, and (vi) oyster bed. The epibenthic community assemblage of the mud habitat was significantly different to that of sand, which in turn differed from the structured habitats of coral reef, mixed reef and oyster bed, with the macroalgal assemblage having similarities to both sand and the other structured habitats. Fish assemblages derived from video data did not differ between habitats, although certain species were only associated with particular habitats. Epibenthic diversity indices were significantly lower in mud, sand and macro-algal habitats, with no differences recorded for fish diversity. Soft sediment grab samples indicated that mud habitats had the highest benthic diversity, with Shannon-Weiner values of >4 , and were more diverse than sand with values of 3.3. The study demonstrates high biodiversity in benthic habitats in the central and southwestern Gulf, which may in part be due to the absence of trawling activity in Qatari waters. There is a strong influence of depth on benthic habitat type, so that depth can be used to predict habitat distribution with a high level of accuracy. The presence of outcrops of hard substrata creates a mosaic of patchy shallow structured benthic habitat across extensive areas of the offshore seabed. Such heterogeneity, and the association of commercially exploited fish species with specific habitats, indicates that this region is well suited to a spatial approach to fisheries management.

Keywords: Arabian Gulf, benthos, biodiversity, fish density, habitat, ROPME Sea Area, spatial management.

Introduction

Marine habitats in the Gulf, in the ROPME Sea Area, face significant anthropogenic pressure resulting from oil and gas extraction,

coastal reclamation, fishing and pollution (Sale *et al.*, 2011; Naser, 2014). These pressures occur against a background of an increasing frequency of extreme temperature events (Riegl *et al.*,

2011). Understanding and managing the impacts of these pressures on marine habitats are severely compromised by lack of data on the distribution and extent of habitats that exist in the Gulf, (UNEP, 2009). Since the introduction of the Convention on Biological Diversity countries became legally obliged to develop a strategy for the conservation and sustainable use of biological diversity, and the importance of marine and coastal biodiversity was recognised early on at the first Conference of Parties (COP 1). All the Gulf States have either ratified or acceded to the CBD and have either produced a National Biodiversity Strategy and Action Plan (NBSAP) or are implementing measures to satisfy obligations to the CBD (CBD, 2016). NBSAPs consist of strategic goals that include setting up of protected areas, protecting and conserving marine and coastal resources, enforcing environmental legislation and supporting scientific research that can better inform decision makers (CBD, 2016). In 2006 a target of 10% of all ecological regions were to be protected by 2012; this was later revised at the COP 10 increasing the terrestrial target to 17% but the marine target was maintained at 10% by 2020 (CBD, 2010). However, in order to achieve these targets countries will need to understand better the inventory of different benthic species and habitats that occur in their national waters within the EEZ.

The unique habitats that exist in the Gulf are a consequence of the sea's isolation and extreme environment due to the restricted water exchange with the Gulf of Oman, combined with high evaporation ($>2,000 \text{ mm yr}^{-1}$) (Hunter, 1983) and low freshwater input that results in great seasonal water temperature variation (10–40 °C) and high salinity (36–60 ppt) (Sheppard *et al.*, 1992; Al-Maslamani *et al.*, 2007). The low energy environment of the Gulf means that the dominant benthic habitat is mud which can stretch from intertidal salt marshes to the deepest waters at around 90 m, with coarser sediment occurring in areas with greater hydrodynamic energy (Sheppard *et al.*, 2010). Intertidally and in shallow waters, these muddy habitats can be extremely productive due to the formation of dense algal mats dominated by cyanobacteria and diatoms that support coastal food webs and commercially important species including both fish and shrimp (Al-Zaidan *et al.*, 2006). Seagrass beds are another highly productive habitat with above-ground biomass of as much as 900 g DW m^{-2} on the east coast of Qatar (Walton *et al.*, 2016), more than 10 times than reported for this species from elsewhere in the world (Duarte and Chiscano, 1999). Macroalgae can form dense beds where hard substrata provide point of attachment, and form an important component of the detrital loop (Sheppard *et al.*, 2010). The apparent high productivity in these habitats is consistent with the idea of Jones *et al.* (2002) that shallower subtidal and intertidal areas in the Gulf are more productive than deeper offshore waters, with recent studies suggesting that much inshore fauna appears to be nutritionally reliant on these habitats (Al-Zaidan *et al.*, 2006; Al-Maslamani *et al.*, 2007; Walton *et al.*, 2014). Among the predominant soft sediments found offshore, shallower limestone outcrops also occur, composed of ancient limestone domes or more recent diagenetic limestone, that are colonised by benthic epifauna including corals and oysters (Riegl, 1999, Sheppard *et al.*, 2010). While lacking the structural complexity of tropical reefs, these oyster and coral habitats are important in terms of both faunal biodiversity and biomass, and are one of the most productive habitats in the Gulf (Sheppard *et al.*, 1992). In areas closer to the coast, patchy corals may colonise areas of hard substrate with better developed coral reefs occurring further offshore (Sheppard *et al.*, 2010). Pearl oyster beds

(*Pinctada radiata* oyster beds occur mostly on hard substrata at depths of between 12 and 16 m (Al-Khayat and Al-Ansi, 2008) and sometimes within patchy coral areas (Smyth *et al.*, 2016). The literature records more than 22 named offshore limestone outcrops with coral rich areas and oyster beds in Qatar waters alone and preliminary studies report they are associated with at least 158 and 189 species, respectively, with many of the species common to both habitats (Al-Ansi and Al-Khayat, 1999; Al-Khayat and Al-Ansi, 2008). These shallow offshore hard substrate habitats and especially oyster beds are targeted by trap fishermen (Al-Maslamani *et al.*, in prep), but to date there are no reports of linkages between habitats in the Gulf and associated fisheries.

While there is a body of knowledge about coastal and shallow (<6 m) water habitats in the Gulf (Al-Zaidan *et al.*, 2006; Al-Maslamani *et al.*, 2007; Walton *et al.*, 2014), they are reported to only occupy <20% of the Gulf (Sheppard *et al.*, 2010). The majority of the $35,000 \text{ km}^2$ of seabed that comprises the Qatar EEZ is comprised of soft sediment habitats in relatively deeper waters (CBD, 2016), which are less well studied and little is known about their importance to associated epibenthic and demersal communities.

Many of the species that occur in benthic habitats in the Gulf are already at the edge of their distribution and are subject to significant environmental stress, and this is compounded by the range of users competing for access to coastal and offshore marine space and resources (Sale *et al.*, 2011), so that additional pressure from climate change or anthropogenic activities resulting from the rapid coastal development and petrochemical industries can have significant impacts. A number of studies have reported that species diversity in the Gulf is impoverished (e.g. Price, 1982; Sheppard *et al.*, 1992; Sheppard *et al.*, 2010), although this is not a universal view (Price and Izsak 2005). Commercial fisheries in Qatar are largely artisanal, and the majority of landings are demersal species caught with traditional baited and non-baited traps (gargoor) (FAO, 2003). Bottom trawling has been banned since 1992 (Decree 86 issued by Ministry of Municipal Affairs and Agriculture in 1992) and prior to that a closed season (1st Feb to 30th June) had been in effect since 1981 (Decree 30 issued by the Council of Ministers, 1981). Bottom trawling is now recognized as one of the most destructive methods of fishing, modifying the seabed morphology and composition of the benthic community (Kaiser *et al.*, 2002) and decreasing both benthic biodiversity and biomass (Hinz *et al.*, 2009). Thus, benthic habitats in Qatar would be expected to be relatively intact compared with neighbouring countries where trawling has continued. This study offers an excellent opportunity to examine the heterogeneity and diversity of offshore benthic habitats utilised by commercial fish species and to inform potential future establishment of ecosystem-based approaches to spatial management of fisheries (Norse, 2010).

Qatar's national waters contain habitats representative of those found in the central and southwestern Gulf, with the same range of environmental stressors as found elsewhere in the region. The present study characterises the varied habitats that are present in the offshore waters of Qatar, and tests the hypothesis that these habitats and associated communities are depth dependent. The mapping of benthic habitats in the present study enables an initial quantitative analysis of the commercially important fish species that are associated with each habitat, as a first step towards an ecosystem based approach to the management of habitats and associated fisheries.

Table 1. Habitat classification and selection criteria based on the number of benthic biota (macroalgae, oyster and coral) and percentage cover of sediment [*sediment classification validated by grain size analysis (see results section)].

Criteria/habitat	Macroalgae	Oyster	Coral	Sand	Mud	Sediment based on camera images*
Macro algal	≥15	Absent	<10			–
Oyster bed	≤25	>10	<10			–
Coral reef	<10	<10	>10			–
Mixed reef	>10	Present	>10			–
Sand	<10	<10	<10	>80%	<20%	Some grains >1 mm visible
Mud	<10	<10	<10	<20%	>80%	No grains >1 mm visible

Methods

Description of habitats

Forty eight stations were sampled at depths between 12 and 52 m using the Qatar University research vessel, RV Janan. At each station HD video and photograph images were collected using a towed camera sledge, with a forward facing GoPro camera angled at 45° and a downward facing digital SLR which was set to take a photo every 20 s. Parallel laser pointers were attached to the front of the frame to calibrate the scale of images.

Camera sledge tows had a duration of 20–30 min, with the exact length determined retrospectively from GPS, speed of the research vessel and total bottom time. Each tow was divided into sections for the analysis of fauna. Each of these sections was 0.178 m × 25 m which gave a sample area of 4.45 m² in total for each separate section.

For sites with soft sediments, quantitative infauna samples were also collected using three replicate 0.1 m² Day grabs, at each station. Each sample was mixed and divided in two. Half was processed for grain size analysis and the other preserved in 70% ethanol for the identification of benthic assemblage. Samples were washed over 0.5 mm mesh, and retained sediments were then examined under a stereo microscope and all organisms identified to species level when possible.

Video habitat classification

Following Ierodiaconou *et al.* (2007) the video images (digital) were initially viewed to determine the types of substrata and dominant biota present. Using these preliminary observations of the image data, a habitat classification was devised using a combination of the percentage cover of a sediment type and counts of dominant epibenthic taxa present in each quadrat using decision rules (Table 1).

Species biodiversity

All epibenthic organisms and fish were recorded and identified from the video footage to either family or species level (S.I. 1). If species level identification could not be achieved, taxa within a genus or family were separated according to morphological difference following the CATAMI classification scheme of Althaus *et al.* (2015), with the image of each taxa archived for comparison with subsequent video footage. Macroalgae were included as pooled classification as it was not possible to separate them due to the variable lighting and image resolution.

Data for fish assemblages and epibenthic communities were analysed by combining both univariate and multivariate analyses, using statistical software PRIMER-E 6 with PERMANOVA extension and SPSS (Clarke and Gorley, 2006, SPSS, 2013). A fourth root transformation was applied to the abundance data collected

Table 2. Depth classes and their associated depth range.

Depth category	Depth range (m)
1	10–20
2	21–30
3	31–40
4	41–50
5	≥51

to improve the data distribution throughout the analysis (Quinn and Keough, 2002). In-fauna data was only available from the two soft sediment habitats and was therefore excluded from ANOSIM and SIMPER analysis.

Multivariate one-way analysis of similarities (ANOSIM) tested the hypothesis that there are significant differences in epibenthic community assemblages between habitats, and these change with increasing depth. Multidimensional scaling (MDS) plots allowed a comparison of computed Bray–Curtis similarities between all sites. SIMPER analysis (analysis of contribution to similarity) was used to identify individual epibenthic species that contributed most to dissimilarities between habitats grouped according to the results from the ANOSIM analysis.

Species diversity indices (Species richness (S'), Pielou's evenness (J'), Margalef Diversity Index (d'), and Shannon–Wiener ($S-W$) Diversity Index (H')) were obtained using the DIVERSE function in PRIMER using the lowest taxonomic groupings the epibenthic fauna from the video footage (SI 1) and the infauna identified from three replicate grab samples. These indices were tested for normality using a Kolmogorov–Smirnov test before being subjected to either a one-way-ANOVA ($P < 0.05$) or Kruskal–Wallis test ($P < 0.05$) to determine if the indices were significantly different. If significance differences were found, a post-hoc analysis (Tukey or Mann–Whitney U) was implemented to determine which groups were significantly different.

Depth analysis

Once each transect was associated with a particular habitat type and depth category (Table 2) a Chi-Square test for association was applied to determine the relationship between depth and habitat type.

Habitat distribution

A multinomial probit analysis was used to determine the depth distribution for each of the habitats. Preliminary inspection of results indicated that coral reef, oyster bed, macroalgae, and mixed reef habitat depth ranges overlapped, hence thereafter they were combined in into a single category of “structured habitat.”




Figure 1. Location of sample stations (circle) and bottom trawl surveys (star). Contour lines show 20 m depth increments. Inset map shows the Gulf and survey area off Qatar.

The probit analysis showed transition zones at intermediate depths between the structured/sand habitats and the sand/mud habitats, so these were added as classification categories. Once the habitat relationship with depth was determined, a map of the possible distribution of these habitats was created. The map was created by using ArcGIS (10.2.1) (ESRI, 2011), with a bathymetric chart obtained from GEBCO and by using the appropriate depth ranges associated with each habitat (GEBCO, 2015). The accuracy of the predicted habitat map was determined by the use of a linear regression t-test, which tested the observed habitat types against the predicted habitat types.

Bottom trawling for commercial fish species

Bottom trawls were carried out as part of a wider fisheries survey that was performed by Department of Fisheries, Ministry of Municipality and Environment (Qatar) between 2008 and 2012. Of these, 16 stations to the north and east of Qatar were stratified according to depth (Figure 1). Five trawls were performed at each of 16 stations at mean (\pm SD) depths of between 9.3 ± 1.5 and 67.6 ± 5.0 m (see Supplement S1). Bottom trawls were carried out at 3 knots for 30 mins using a net with headline height of 1 m, a spread between doors of 19 m, with a 54 mm mesh codend and wing panels. The catch was sorted into commercially important species counted and weighed wet on motion compensated balances accurate to 1 g. Biomass data for each species was expressed as catch per unit area based on swept area using GPS positions for the start and end of trawl.

Results

Epibenthic species compositions in each habitat

Video footage indicated macroalgae were present in all habitats, with the greatest frequency occurring in the “macroalgal” and “mixed reef” habitat categories (Table 3). Similarly, corals appear with the greatest frequency in the coral reef and mixed reef habitats. Higher taxonomic level identification (presented in Supplement S1) shows that sea pens, polychaetes, and Actiniaria

were found almost exclusively in the sand and mud habitats. All other genera/species recorded during the camera sledge tows were not habitat specific and were recorded in almost all habitats (Table 3).

MDS analysis indicated that the benthic assemblages found over the sampled habitats clustered into three main groups: mud, sand and the photic-zone structured habitats (SH) (the latter comprised of macroalgal, coral reef, mixed reef, and oyster bed) (Figure 2). There were significant differences between the species groupings found across these habitats (ANOSIM, $r=0.636$, $P \leq 0.001$). A pairwise comparison between all habitats confirmed that benthic assemblages were significantly different (Table 4).

SIMPER analysis of the epibenthic community assemblages, showed macroalgae, ascidians (*Phallusia nigra*), the black long spine urchin *Diadema setosum*, and sea pens and sea whips are the main taxa responsible for dissimilarity between the habitats. The greatest dissimilarities were seen between the mud and structured habitats (90.32%) that were separated by the greatest depth difference (Table 5).

Infaunal community

Grain size analysis of the soft sediment grab samples revealed significantly more sand and less clay in the shallower “sand” stations than the deeper “mud” stations ($t=28.3$, $P < 0.001$, and $t=6.3$, $P < 0.001$, respectively). From the grab samples 603 species were identified and more than 13 000 individuals were counted. The composition of the benthic infauna indicated that annelids and arthropods were the two most important phyla in terms of abundance (Table 6). These phyla, together with other abundant in-fauna such as nemertean, sipunculids and molluscs are not well captured in the video footage. Significantly greater abundances of annelids, arthropods, nemerteans, and sipunculids were recorded in the shallower sand habitats than in the mud habitats.

Fish species compositions in each habitat (camera data)

In relation to the mobile fish species, *Valenciennea persica* was recorded only in the macroalgae habitat. *Nemipterus* spp., *Lepidotrigla bispinosa*, *Selaroides leptolepis*, and *Upeneus* spp. were found only in the mud habitat. *Pseudochromis aldbaeraensis* was recorded exclusively in coral reef habitats and *Dussumieria* spp., *Epinephelinae* spp., and *Parupeneus margaritatus* were recorded only in sand habitats. Gobiidae in burrows were observed in all habitats. All other fish species recorded during the camera tows were not limited to a single habitat (Table 7).


In order to improve the statistical power coral reef, mixed reef, oyster bed and macroalgae habitats were pooled. The mixed distribution in MDS ordination plots suggested fish species were not discriminated by habitat type (Figure 3) and was confirmed by ANOSIM which showed no significant differences between the assemblages of fish in the three habitats (structured, sand and mud, ANOSIM, $r=0.173$, $P=0.074$).

Community diversity

Diversity indices, using the lowest identified taxonomic levels identified from the video footage, indicated that mixed reef habitats had the highest epibenthic diversity compared with the other sampled habitats (Table 8). Sand and mud habitats had the lowest diversities for all indices. Habitats (excluding the grouped structured habitat) were found to have significantly different species

Table 3. The percentage occurrence of epibenthic taxa within a 4.45 m² quadrat in each habitat type (%) identified from camera images.

Taxa	Oyster Bed	Mixed Reef	Coral Reef	Macroalgae	Sand	Mud
Acroporidae	37.5	7.7	26.7	8.3	0	0
Actiniaria	0	0	0	0	0.8	11.7
Ascidiae	100	76.9	60.0	58.3	13.1	1.0
Chalinidae	0	30.8	13.3	0	0.4	0
Clypeasteroidea	37.5	23.1	6.7	8.3	5.6	0.5
Demospongiae	75.0	7.7	40.0	16.7	9.1	3.1
Dendrophylliidae	87.5	100	60.0	25.0	0.4	0
Diadematidae	62.5	61.5	100	0	1.6	0.5
Holothuriidae	12.5	0	33.3	8.3	0.4	0
Macroalgae	100	100	26.7	100	38.5	4.1
Mussidae	25.0	69.2	100	0	0.8	0
Ophiasteridae	25.0	61.5	13.3	0	0.8	0.5
Pennatulacea	50.0	0	6.7	25.0	11.9	65.3
Poritidae	62.5	46.2	86.7	0	1.2	0
Pteriidae	100	69.2	20.0	16.7	0.4	0
Sabellidae	0	0	0	0	4.0	1.0
Siderastreidae	12.5	46.2	40.0	0	0.4	0

**Figure 2.** Multi-dimensional scaling plot of the similarity in the community composition of epibenthic species (average density per 4.25 m²) for mud, sand, and structured habitats (SH) sampled by towed camera survey.**Table 4.** Pairwise comparison of benthic community composition in the sampled habitats.

Habitats	R Statistic	P Value
Mud and sand	0.616	0.001
Mud and structured	0.937	0.001
Sand and structured	0.378	0.001

richness ($X^2_{(5)} = 19.35$, $P = 0.002$), total individual frequency ($X^2_{(5)} = 16.53$, $P = 0.004$), Margalef diversity index (ANOVA $F_{(5,31)} = 10.47$, $P < 0.001$) and Shannon–Wiener (S–W) diversity index ($X^2_{(5)} = 19.14$, $P = 0.002$), while no significant difference were found using Pielou’s evenness ($X^2_{(5)} = 6.72$, $P = 0.242$). Post-hoc analysis showed that oyster beds and mixed reef had significantly greater Margalef diversity and S–W diversity than mud and sand habitats. Species richness and total individual frequency indices also indicated significantly higher diversity in coral reef compared with mud and sand habitats. Comparison of the

epibenthic diversity in the grouped structured habitat with that of mud and sand indicated significant differences in Margalef diversity index (ANOVA $F_{(2,35)} = 3.497$, $P = 0.041$), S–W diversity index (ANOVA $F_{(2,35)} = 5.359$, $P = 0.009$), species richness ($X^2_{(2)} = 11.81$, $P = 0.004$), total individual frequency ($X^2_{(2)} = 14.13$, $P = 0.001$), while no significant difference were found using Pielou’s evenness (ANOVA $F_{(2,35)} = 0.788$, $P = 0.467$). Post hoc analysis showed that structured habitats had significantly greater Margalef diversity, Shannon–Wiener diversity, total individual frequency and species richness than sand and mud habitats.

Analyses of the Day grab sampled soft sediment communities indicated benthic diversity to be much greater than that captured in the camera data (Table 9). Grab samples indicated S–W diversity was higher in mud habitats ($T\text{-test}_{(17)} = 6.44$, $P < 0.001$), Pielou’s evenness was also higher in mud habitats (Kruskal–Wallis $X^2_{(1)} = 19.46$, $P = 0.242$), but total individual frequency was higher in sand habitats ($X^2_{(1)} = 15.98$, $P < 0.001$). Both S–W diversity and Pielou’s evenness were significantly positively

Table 5. Results of SIMPER analysis showing the top three taxa that differed between habitat classifications.

Habitat (average dissimilarity)	Species	Average dissimilarity	Contribution to dissimilarity (%)	Habitat where in greatest abundance
Sand and mud (78.64%)	Sea pens	11.28	14.34	Mud
	Macroalgae	9.81	12.47	Sand
	Sea whips	6.68	8.50	Mud
Sand and structured habitat (73.78%)	<i>P. nigra</i>	6.16	8.36	Structured
	Macroalgae	5.71	7.73	Structured
	<i>D. setosum</i>	5.62	7.62	Structured
Mud and structured habitat (90.32%)	Macroalgae	8.97	9.93	Structured
	Sea pens	8.18	9.06	Mud
	<i>P. nigra</i>	7.00	7.75	Structured

Average dissimilarity in abundance and contribution to the overall dissimilarity is given.

Table 6. Mean abundances (inds. m⁻²) of phyla recorded in grab samples of the infauna, (division level in the case of Rhodophyta).

Classification	Mud	Sand
Rhodophyta	0	0.33
Sarcomastigophora	12.0	11.56
Porifera	0	0.11
Cnidaria	0.657	4.33
Nemertea	0	0.33
Nematoda	5.43	26.22
Sipuncula	1.14	26.67
Platyhelminthes	0	0.56
Annelida	163.986	386.33
Arthropoda	70	82.56
Mollusca	35.986	33.22
Echinodermata	1.14	5.33
Brachiopoda	0	0.33
Ectoprocta	3.71	2.89
Chordata	0	1.67

correlated with increasing depth of habitat (Pearson correlation $r = 0.75$, $P < 0.001$ and $r = 0.79$, $P < 0.001$, respectively). No significant differences were detected in species richness and Margalef diversity between mud and sand habitats.

Video data showed greater fish species richness associated with mud and macro-algal habitats as well as total individual frequency, Margalef diversity index, and S–W diversity. Pielou's evenness values were relatively constant for fish species across all habitats (0.99) with the exception of mixed Reef which was lower (0.95, Table 10). However, DIVERSE analysis of the fish community showed no significant differences in the diversity indices for the sampled habitat categories for fish species richness ($X^2_{(2)} = 1.858$, $P = 0.395$), total individual frequency ($X^2_{(2)} = 1.623$, $P = 0.444$), Pielou's evenness (ANOVA $F_{(2,19)} = 0.087$, $P = 0.917$), S–W diversity ($X^2_{(2)} = 0.386$, $P = 0.386$), and Margalef diversity index ($X^2_{(2)} = 1.934$, $P = 0.380$).

Habitats with depth

The most abundant habitats in the 10–20 m depth range were the structured habitats (coral/mixed reef, oyster bed, and macroalgae) which represented 53.3% of the observed habitat types. From 21 to 30 m, only one structured habitat (macroalgae) was present, while the proportion of sand and mud habitats increased. From 31 to 40 m, only mud and sand habitats were found, and below 40 m mud was the only habitat observed. A chi-square test for association determined that there was a significant relationship

between habitats observed and depth ($X^2_{(20)} = 616.156$, $P \leq 0.001$). There was a strong association between depth and habitat type (Cramér's $V = 0.564$, $P \leq 0.001$) (Figure 4).

Prediction of spatial distribution of habitats

Using the close relationship between habitat and depth data, multinomial probit analysis determined the depth range at which each of the habitats were likely to be present as structured <14 m, structured to sand 14–19 m, sand 20–27 m, sand to mud 26–35 m, and mud >36 m. Transitional habitats, e.g., sand to mud were composed of a mosaic of sand and mud areas. These groupings were used to predict spatial distributions for each habitat (Figure 5). A linear regression t-test indicated there was no difference between the observed and predicted habitats at each site ($t(88) = 1.91$, $P = 0.06$), with the habitat prediction map achieving an accuracy of 93.48%.

Commercial fisheries trawl survey

The survey trawls covered 0.003–0.004% of the trawlable area within the Qatar EEZ, with individual trawls covering approximately 0.05 km². Resulting mean catch rates ranged between 547 and 740 kg km⁻², of which less than half the biomass was due to the 12 most exploited commercial species. Fish catches over the habitat classifications indicates that some species such as the lethrinids and the sparids, with the exception of *Agyrops spinifer*, were more abundant over the shallow structured habitats, while the groupers and *A. spinifer* were found over the deeper sand to mud habitats (Figure 6). The transitional habitats (structured-sand, sand-mud) had the highest mean (\pm SE) fish biomass and resulted in the highest catches of commercial species per trawl (structured—58.9 \pm 42.4 kg km⁻², structured to sand—171.3 \pm 39.0 kg km⁻², sand—80.3 \pm 30.5 kg km⁻², sand to mud—121.1 \pm 43.6 kg km⁻², and mud—34.4 kg \pm 10.2 km⁻²). However, analysis of variance suggested there was no significant difference ($F = 2.25$, $P = 0.072$) between the log transformed catch rates from the different habitats.

Discussion

In the offshore waters west of Qatar (10–70 m) six habitat types were identified using the guidelines presented in Ierodiaconou *et al.* (2011): two soft sediment habitats of mud and sand and four structured habitats: coral reef, mixed reef, macro-algal and oyster bed. Sea bed video footage was used as a cost-effective method to identify the range of epibenthic species present over each of these habitats. While identification was not always possible to species level this is not problematic as the use of coarser

Table 7. The percentage occurrence of the most abundant fish families found in each habitat type within a 4.45 m² quadrant (%) and with the coral reef, mixed reef, oyster bed, and macroalgae grouped into the structured habitats.

Taxa	Coral Reef	Oyster Bed	Mixed Reef	Macroalgae	Structured = CR&MR&OB&MA	Sand	Mud
Carangidae	–	–	–	–	0	0.4	1.0
Carcharhinidae	–	–	–	–	0	0.4	–
Clupeidae	–	–	–	–	0	–	0.5
Dussumieriidae	–	–	–	–	0	0.8	–
Gobiidae	–	12.5	–	25.0	9.1	6.3	14.8
Lutjanidae	–	–	–	16.7	4.5	0.4	2.6
Mullidae	–	12.5	–	–	2.3	6.3	1.0
Paralichthyidae	–	–	–	–	0	0.4	0.5
Pomacanthidae	9.1	12.5	–	8.3	6.8	0.4	–
Pseudochromidae	9.1	–	–	–	2.3	–	–
Serranidae	9.1	12.5	–	–	4.5	0.8	1.0
Sillaginidae	–	–	–	8.3	2.3	4.4	9.7
Sphraenidae	–	–	–	–	0	–	2.0
Synodontidae	–	–	–	–	0	–	4.6
Triglidae	–	–	–	–	0	–	4.6
Unidentified.	–	–	7.7	–	2.3	2.0	4.1




Figure 3. Multi-dimensional scaling plot based on fish species average density per 4.25 m² for mud, sand, and structured habitats (SH) sampled by towed video. The plot shows similarities between sites in regards to their fish community composition in each habitat.

Table 8. Mean values (\pm SE) for the epibenthic community in each habitat sampled by video surveys of species richness (S), total individual frequency (N), Margalef diversity index (d'), Pielou's evenness (J'), and Shannon–Weiner diversity index ($H'(\log_e)$).

Habitat	S	N	d'	J'	$H'(\log_e)$
SH-Macroalgae	4.00 (\pm 2.00)	7.09 (\pm 3.87)	1.47 (\pm 0.62)	0.92 (\pm 0.03)	1.16 (\pm 0.54)
SH-Coral reef	9.50 (\pm 2.50)	19.14 (\pm 3.99)	2.85 (\pm 0.65)	0.96 (\pm 0.02)	2.13 (\pm 0.31)
SH-Oyster bed	9.67 (\pm 1.33)	17.28 (\pm 2.74)	3.04 (\pm 0.33)	0.98 (\pm 0.00)	2.19 (\pm 0.15)
SH-Mixed reef	11.00 (\pm 2.00)	24.27 (\pm 5.89)	3.14 (\pm 0.39)	0.95 (\pm 0.02)	2.27 (\pm 0.22)
SH	10.33 (\pm 1.29)	367.67 (\pm 97.22)	1.64 (\pm 0.21)	0.57 (\pm 0.06)	1.34 (\pm 0.19)
Sand	3.59 (\pm 0.36)	5.54 (\pm 0.74)	1.52 (\pm 0.13)	0.96 (\pm 0.01)	1.15 (\pm 0.10)
Mud	2.73 (\pm 0.38)	4.46 (\pm 0.62)	1.14 (\pm 0.15)	0.94 (\pm 0.01)	0.88 (\pm 0.11)

taxonomic classification when combined with morphological differences correlates well with actual species diversity (Bell and Barnes, 2001; Althaus et al., 2015; Monk et al., 2016). Epibenthic species assemblages differed between habitats, with certain organisms being characteristic of a habitat such as sea pens in mud


habitats. While sand habitats were noticeably impoverished in terms of epibenthos, low densities of all the species recorded in the other surveyed habitats also occurred here. This is in notable contrast to previous work that found higher diversity in sand compared with mud subtidal habitats (Sanders, 1968). Although

Table 9. Mean values (\pm SE) for the infauna community in each habitat sampled by grab of species richness (S), total individual frequency (N), Margalef diversity index (d'), Pielou's evenness (J'), and Shannon–Weiner diversity index (H' log_e).

Habitat	S	N	d'	J'	H' (log _e)
Sand	107.36(\pm 5.13)	618.79 (\pm 60.02)	16.63 (\pm 0.60)	0.72 (\pm 0.02)	3.34 (\pm 0.09)
Mud	96.21(\pm 3.17)	309.93 (\pm 15.95)	16.62 (\pm 0.46)	0.88 (\pm 0.01)	4.00 (\pm 0.04)


Table 10. Mean values of species richness (S), total individual frequency (N), Margalef diversity index (d'), Pielou's evenness (J'), and Shannon–Weiner diversity index (H' log_e) for fish community in each habitat sampled by video surveys (\pm standard error).

Habitat	S	N	d'	J'	H' (log _e)
SH-Macroalgal	5.00 (\pm 0.00)	5.38 (\pm 0.00)	2.38 (\pm 0.00)	1.00 (\pm 0.00)	1.61 (\pm 0.00)
SH-Oyster Bed	4.00 (\pm 0.00)	4.83 (\pm 0.00)	1.91 (\pm 0.00)	0.99 (\pm 0.00)	1.37 (\pm 0.00)
SH-Coral Reef	3.00 (\pm 0.00)	3.00 (\pm 0.00)	1.82 (\pm 0.00)	1.00 (\pm 0.00)	1.10 (\pm 0.00)
SH-Mixed Reef	2.00 (\pm 0.00)	2.68 (\pm 0.00)	1.01 (\pm 0.00)	0.95 (\pm 0.00)	0.66 (\pm 0.00)
SH	3.50 (\pm 0.65)	3.97(\pm 0.67)	1.78 (\pm 0.28)	0.98 (\pm 0.01)	1.18 (\pm 0.20)
Sand	3.67 (\pm 0.91)	4.49 (\pm 1.18)	1.73 (\pm 0.27)	0.99 (\pm 0.20)	1.10 (\pm 0.02)
Mud	5.00 (\pm 1.13)	6.21 (\pm 1.40)	2.14 (\pm 0.32)	0.99 (\pm 0.19)	1.42 (\pm 0.02)

**Figure 4.** Relationship between depth and percentage frequency of benthic habitat categories in the offshore waters of Qatar. SH = structured habitat.

more recently it has been suggested that grain size alone is unlikely to result in the diversity differences reported as many other variables are co-correlated (Snelgrove and Butman, 1995).

The differences in species assemblages found between the complex limestone-outcrop structured habitats of oyster bed, mixed reef and coral reef and the homogeneous soft-sediment sand and mud habitats were further supported by the diversity indices. These indices also indicated significantly greater epibenthic diversity over the more complex habitats compared to those more homogeneous sand and mud habitats. Previous studies have reported low diversity in the Gulf (Price, 1982; Sheppard *et al.*, 1992) due to the stress caused by the harsh environmental conditions of extreme sea temperatures (10–40 °C) and high salinities (36–60 ppt) (Sheppard *et al.*, 1992; Al-Maslamani *et al.*, 2007), combined with the geographic semi-isolation of the Gulf due to the restricted water exchange through the Strait of Hormuz (Wilson *et al.*, 2002). Comparable epibenthic diversity studies using a towed camera system are rare, but one study in Cardigan Bay Special Area of Conservation (SAC) on the Welsh coast found very similar S–W diversity levels of between 1.2 and 2.6 over sand and gravel at depths of 3–12 m (Sciberras *et al.*, 2013)

**Figure 5.** ArcGIS map predicting habitats present in Qatari Waters to the North-East of Doha. Contour lines are depth increments of 20 m.

compared with the current study in Qatar waters where S–W diversity varied between 0.9 and 2.3. Similar levels of S–W diversity were also reported using towed cameras on the Greenland continental shelf on both hard and soft substrates, of 0.1–2.5 (Yesson *et al.*, 2015) and using autonomous underwater vehicle reef and soft sediments in Tasmania of 0.13 and 1.98 (Monk *et al.*, 2016). Trawling for benthic epifauna from the organically polluted Hong Kong harbour to the less polluted outside channel, indicated the epifaunal S–W diversity indices increased with distance from the harbour from 0.7 to 2.1 (Wu, 1982). Overall, these comparisons suggest that the epibenthic diversity in the central Gulf is not impoverished relative to comparable depths in other biogeographic regions. This may well reflect intactness of offshore habitats as a result of the bottom trawling ban implemented by Qatar




Figure 6. Mean trawl catch rates and standard error (kg km^{-2}) for the 12 most commercially important species.

in 1994 (UN, 1997), as especially in the deeper areas, sea pens and sea whips were commonly recorded in almost 60% of the 4.45 m² quadrants in mud habitats. However, although some fish species were only found in certain habitats, neither fish diversity nor fish species composition based on camera data differed significantly between habitats. Slightly higher infauna S–W diversity (converted to H’log_e) has been reported in similar muddy habitats and depths in the Indian Ocean off Madagascar (4.4), with lower diversity in the Arabian Sea (2.1 off Mumbai and 1.8 off Kerala) although these might be the result of an annual hypoxia event (Sanders, 1968). However, the deeper muddy areas sampled in the current Gulf study are also thought to experience a similar

hypoxia event during the summer (Al-Ansari et al., 2015). The first quantitative study on trawling impacts, found that increasing bottom trawling activity from 1.3 to 18.2 times a year decreased infauna abundance by 72%, biomass by 77% and species richness by 40% (Hinz et al., 2009). Similar inference have been made from a study on Georges Bank with areas disturbed by dredges/trawling having a lower S–W diversity (2.7) than undisturbed areas (3.2) (Collie et al., 1997). The benthic diversity of the soft sediments sampled with grabs in the current study was surprisingly rich, with higher S–W diversity indices in the muddy furthest offshore area of 4 and lower values in the closer sandier habitat of 3.34 where benthic disturbance as a result of trap-based

fishing activity is thought to be higher (M. Abdallah, pers. comm.).

Benthic habitat type in the Gulf waters around Qatar were shown to be strongly correlated with depth. In the offshore shallower waters (10–20 m) four structured habitats on limestone outcrops were sampled, coral reef, oyster beds, macroalgae, and mixed reef; these transitioned to sand and then to mud habitats with increasing depth. Multinomial probit analysis resolved the depth transitions between habitats and enabled the production of a predictive habitat map. In the Gulf the photic zone only extends down to 6–15 m (Sheppard *et al.*, 2010), which is approximately the same depth as the structured habitats. The depth of the photic zone is limited by the resuspension and addition of sediment that results from strong winter North-Westerly winds (“shamal”) (Alsharhan and Kendall, 2003) which increase turbidity in the water column reducing light penetration. The combination of low light levels and high sedimentation would lead to increased stress levels for photic organisms close to their photosynthetic compensation depth. The frequency of sedimentation events combined with fluctuations in temperature and salinity are likely responsible for the reduced number of coral species in the area (Morelock *et al.*, 1979). The lack of hard substrate in deeper waters further restricts the distribution of photic organisms to the shallow waters over the limestone outcrops (“hairat”), which are geological rather than biogenic features. Oysters beds (*Pinctada radiata*), depend on phytoplankton and the availability of a suitable settlement substrate and thus appear to be restricted to the photic zone. The resuspension of sediment caused by increased water turbulence during storms and shamals facilitates the movement of fine particles from shallower depths to the deeper low energy waters found offshore (Monroe and Wicander, 2011).

Comparing seabed habitats in Qatar, epibenthic species richness was significantly higher in coral reef, mixed reef and oyster beds, compared with sand and mud habitats. However, the compound diversity indices only found oyster and mixed reef to have significantly higher epibenthic diversity, and none of the habitats had significantly different fish diversity. This is consistent with previous wider regional studies of fish assemblages, which report lower coral-associated fish diversity and biomass within the Gulf compared to the neighbouring waters of the Gulf of Oman and Arabian Sea, and is attributed to the limited extent and impoverished diversity of corals (Burt *et al.*, 2011; Sale *et al.*, 2011). Fish S–W diversity in Qatari waters was found to vary between 0.7 and 1.6. Fish assemblage S–W diversity recorded by divers on coral reefs in Cape Verde varied between means of 1.8 and 2.6 (Santos *et al.*, 2013), and 0.7–4.9 over coral reefs in the Yucatan (Arias-González *et al.*, 2012). Lower diversity was observed in the current study in Qatar, but this was achieved using benthic-focused towed cameras where the field of view might not have captured more mobile fish species. However a diver survey carried out in the same area, found similar S–W diversities of 1.35 and 1.89 over sand and coral reef habitats, respectively (Hayes, 2015, Egerton *et al.*, in prep). Trawl catches of the commercial species indicate that the structured habitats are important for some species such as the lethrinids and sparids, but these species are also found in significant numbers in deeper habitats. Moreover, deeper habitats of >27 m that included the sand-mud and mud habitat classifications are significant areas for other important commercial species such as groupers and *A. spinifer*. It is not known whether the pattern of fish catches observed represents actual habitat use as all trawl surveys were performed during

daylight, and nocturnal habitat use would not be captured in the current study. The lack of difference in fish catches and fish biodiversity recorded between the structurally complex shallow habitats and deeper less complex habitats may also result from the higher fishing pressure that occurs in these shallower areas as these habitats are targeted as traditional fishing grounds. The damage to epifaunal communities in the retrieval of these traps is of concern (Al-Maslamani *et al.*, in prep). Soak times are normally 3–4 days and fishermen do not use marker buoys, only marking the position of their strings of traps with GPS to reduce the risk of theft. The fishermen therefore have to retrieve the traps with a heavy grapple, and this grapple together with the trap recovery causes extensive physical damage to the benthos.

In conclusion, the study demonstrates the strong influence of depth on benthic habitat type, but not fish community composition, so that depth can be used to predict habitat distribution with a high level of accuracy. Although the shallow structured habitats of reef, mixed reef, oyster beds, and macroalgae support higher epibenthic diversity than the deeper habitats (sand, mud) there was no difference in diversity between types of structured habitat. Overall, the presence of outcrops of hard substrata creates a mosaic of patchy shallow structured benthic habitat across extensive areas of the offshore seabed. Such heterogeneity, and the association of commercially exploited fish species with specific habitats, indicates that this region is well suited to a spatial approach to fisheries management of the sort described by Norse (2010). A similar approach has been used in the development of the marine spatial plan of Australia, where distribution of demersal fish and habitats are used to determine benthic bioregions (Commonwealth_of_Australia, 2006) onto which human use maps are overlaid to guide marine spatial zonation (Norse, 2010). In the present study, it was notable that high epibenthic diversity did not translate into high fish diversity and this is supported by the trawl data of the commercial catches that showed trawling over the transitional habitats resulted in the highest catches. Hence, some further work on association of exploited fish species and communities with specific habitats and connectivity between habitats is required, including acoustic studies of fish distribution and abundance over shallow structured habitats (Egerton *et al.*, in press), food web linkages between benthic habitats and demersal fish, and development of better understanding of connectivity through investigation of diurnal, seasonal and ontogenetic patterns of fish movement. Together, this information will give support to potential new management measures, including integration of habitat protection and spatial management of fisheries in the Gulf.

Supplementary data

Supplementary material is available at the *ICESJMS* online version of the manuscript.

Acknowledgements

This research was supported by the project “Identification of Essential Fish Habitat to Achieve An Ecosystem-Based approach to Fisheries Management” funded by the Qatar National Research Foundation (QNRF) NPRP 6-1680-4-027. We would particularly like to thank Jodie Haig, Julia Pantin, Harriet Salomonsen, Gwladys Lambert, Caesar Sorino, Abdulrahman Albinali, Yousif Alkahalaf for participation in survey cruises on the RV Janan, and Ms. Aisha Al-Ashwel, Ms. Najat Al-Omary, and Ms. Noora Ali Alfardi for help with benthic infauna species identification and

Manal Ibrahim, Noora Alshammari for fish laboratory work. We would also like to thank the reviewers and the editor, Dr Norkko, whose thoughtful comments greatly improved the manuscript.

References

- Al-Ansari, E., Rowe, G., Abdel-Moati, M. A. R., Yigiterhan, O., Al-Maslmani, I., Al-Yafei, M. A., Al-Shaikh, I., and Upstill-Goddard, R. 2015. Hypoxia in the central Arabian Gulf Exclusive Economic Zone (EEZ) of Qatar during summer season. *Estuarine Coastal and Shelf Science*, 159: 60–68.
- Al-Khayat, J., and Al-Ansi, M. 2008. Ecological features of oyster beds distribution in Qatari waters, Arabian Gulf. *Asian Journal of Scientific Research*, 1: 544–561.
- Al-Maslmani, I., Le Vay, L., Kennedy, H., and Jones, D. A. 2007. Feeding ecology of the grooved tiger shrimp *Penaeus semisulcatus* De Haan (Decapoda: Penaeidae) in inshore waters of Qatar, Arabian Gulf. *Marine Biology*, 150: 627–637.
- Al-Maslmani, I., Smyth, D., Giraldes, B., Chatting, M., Al-Mohannadi, M., and Le Vay, L. in prep. Is excessive fishing intensity with artisanal gear impeding the recruitment of ecosystem providers onto the historic offshore oyster beds of the western Arabian Gulf?
- Al-Zaidan, A. S. Y., Kennedy, H., Jones, D. A., and Al-Mohanna, S. Y. 2006. Role of microbial mats in Sulaibikhat Bay (Kuwait) mudflat food webs: evidence from delta C-13 analysis. *Marine Ecology-Progress Series*, 308: 27–36.
- Al-Ansi, M. A., and Al-Khayat, J. A. 1999. Preliminary study on coral reef and its associated biota in Qatari waters, Arabian Gulf. *Qatar University Science Journal*, 19: 294–311.
- Alsharhan, A. S., and Kendall, C. G. S. C. 2003. Holocene coastal carbonates and evaporites of the southern Arabian Gulf and their ancient analogues. *Earth-Science Reviews*, 61: 191–243.
- Althaus, F., Hill, N., Ferrari, R., Edwards, L., Przeslawski, R., Schönberg, C. H. L., Stuart-Smith, R., Barrett, N., Edgar, G., Colquhoun, J., Tran, M., Jordan, A., Rees, T., and Gowellett-Holmes, K. 2015. A standardised vocabulary for identifying benthic biota and substrata from underwater imagery: the CATAMI classification scheme. *PLoS One*, 10: e0141039.
- Arias-González, J. E., Acosta-González, G., Membrillo, N., Garza-Pérez, J. R., and Castro-Pérez, J. M. 2012. Predicting spatially explicit coral reef fish abundance, richness and Shannon–Weaver index from habitat characteristics. *Biodiversity and Conservation*, 21: 115–130.
- Bell, J. J., and Barnes, D. K. A. 2001. Sponge morphological diversity: a qualitative predictor of species diversity? *Aquatic Conservation: Marine and Freshwater Ecosystems*, 11: 109–121.
- Burt, J. A., Feary, D. A., Bauman, A. G., Usseglio, P., Cavalcante, G. H., and Sale, P. F. 2011. Biogeographic patterns of reef fish community structure in the northeastern Arabian Peninsula. *ICES Journal of Marine Science: Journal Du Conseil*, 68: 1875–1883.
- CBD. 2010. The strategic plan for biodiversity 2011–2020 and the Aichi Biodiversity Targets. Conference of Parties to the Convention on Biological Diversity at its Tenth Meeting, UNEP/CBD/COP/DEC/X/2, Nagoya, Japan.
- CBD. 2016. CBD-Country Profile-State of Qatar <https://www.cbd.int/countries/profile/?country=qa>.
- Clarke, K. R., and Gorley, R. N. 2006. PRIMER V6: user manual-tutorial. Plymouth Marine Laboratory.
- Collie, J. S., Escanero, G. A., and Valentine, P. C. 1997. Effects of bottom fishing on the benthic megafauna of Georges Bank. *Marine Ecology Progress Series*, 155: 159–172.
- Commonwealth_of_Australia. 2006. A guide to the integrated marine and coastal regionalisation of Australia. Version 4.0. 16
- Duarte, C. M., and Chiscano, C. L. 1999. Seagrass biomass and production: a reassessment. *Aquatic Botany*, 65: 159–174.
- Egerton, J., Al-Ansi, M., Abdallah, M., Walton, M. E. M., Hayes, J., Turner, J. R., Al-Maslmani, I., Mohannadi, M., and Le Vay, L. in prep. Fish association with shallow offshore habitats in the Arabian Gulf.
- ESRI. 2011. ArcGIS Desktop Release 10. Redlands, CA: Environmental Systems Research Institute.
- FAO. 2003. Fishery Country Profile: Qatar. (ftp://ftp.fao.org/FI/DOCUMENT/fcp/en/FL_CP_QA.pdf). FAO, Rome.
- GEBCO. 2015. GEBCO. [Online] <http://gebcocenter.org/>. (last accessed 17 September 2015).
- Hayes, J. 2015. Distribution and species diversity of offshore benthic habitats in the exclusive economic zone of Qatar, Arabian Gulf. MSc Thesis. Ocean sciences, Bangor University. 76 pp.
- Hinz, H., Prieto, V., and Kaiser, M. J. 2009. Trawl disturbance on benthic communities: chronic effects and experimental predictions. *Ecological Applications*, 19: 761–773.
- Hunter, J. 1983. Aspects of the dynamics of the residual circulation of the Arabian Gulf. Pages 31–42 *Coastal oceanography*. Springer. ISBN: 1461566509.
- Ierodiaconou, D., Laurenson, L., Burq, S., and Reston, M. 2007. Marine benthic habitat mapping using Multibeam data, georeferenced video and image classification techniques in Victoria, Australia. *Journal of Spatial Science*, 52: 93–104.
- Ierodiaconou, D., Monk, J., Rattray, A., Laurenson, L., and Versace, V. 2011. Comparison of automated classification techniques for predicting benthic biological communities using hydroacoustics and video observations. *Continental Shelf Research*, 31: S28–S38.
- Jones, D. A., Price, A. R. G., Al-Yamani, F., and Al-Zaidan, A. 2002. Coastal and marine ecology. *In* The Gulf Ecosystem; health and sustainability. pp. 65–103. Ed. by N. Y. Kahn, M. Munwar, A.R.G. Price. *Ecovision World Monograph Series*, Bakhuis, Leiden.
- Kaiser, M. J., Collie, J. S., Hall, S. J., Jennings, S., and Poiner, I. R. 2002. Modification of marine habitats by trawling activities: prognosis and solutions. *Fish and Fisheries*, 3: 114–136.
- Monk, J., Barrett, N. S., Hill, N. A., Lucieir, V. L., Nichol, S. L., Siwabessy, J. P. W., and Williams, S. B. 2016. Outcropping reef ledges drive patterns of epibenthic assemblage diversity on cross-shelf habitats. *Biodiversity and Conservation*, 25: 485–502.
- Monroe, J. S., and Wicander, R. 2011. *The changing earth: exploring geology and evolution*. Cengage Learning. 1133715516.
- Morelock, J., Boulon, K., and Galler, G. 1979. Sediment Stress Coral Reefs. Pages 46–58 *In*: Proceedings, Energy Industry and the Marine Environment in Guayanilla Bay. Centre for Energy and Environmental Research, the University of Puerto Rico.
- Naser, H. A. 2014. Marine ecosystem diversity in the Arabian Gulf: threats and conservation. *In* Biodiversity—The Dynamic Balance of the Planet. pp. 297–328, INTECH; ISBN: 978-953-51-1315-7.
- Norse, E. A. 2010. Ecosystem-based spatial planning and management of marine fisheries: why and how? *Bulletin of Marine Science*, 86: 179–195.
- Price, A. 1982. Comparison between echinoderm faunas of Arabian Gulf, SE Arabia, Red Sea and Gulfs of Aqaba and Suez. *Fauna of Saudi Arabia*, 4: 3–21.
- Price, A. R. G., and Izsak, C. 2005. Is the Arabian Gulf really such a lowspot of biodiversity?: Scaling effects and management implications. *Aquatic Ecosystem Health and Management*, 8: 363–366.
- Quinn, G. P., and Keough, M. J. 2002. *Experimental design and data analysis for biologists*. Cambridge University Press, Cambridge, UK.
- Riegl, B. 1999. Corals in a non-reef setting in the southern Arabian Gulf (Dubai, UAE): fauna and community structure in response to recurring mass mortality. *Coral Reefs*, 18: 63–73.
- Riegl, B. M., Purkis, S. J., Al-Cibahy, A. S., Abdel-Moati, M. A., and Hoegh-Guldberg, O. 2011. Present limits to heat-adaptability in corals and population-level responses to climate extremes. *PLoS One* 6: e24802.

- Sale, P. F., Feary, D. A., Burt, J. A., Bauman, A. G., Cavalcante, G. H., Drouillard, K. G., Kjerfve, B., Marquis, E., Trick, C. G., Usseglio, P., and Van Lavieren, H. 2011. The growing need for sustainable ecological management of marine communities of the Persian Gulf. *Ambio*, 40: 4–17.
- Sanders, H. L. 1968. Marine benthic diversity: a comparative study. *The American Naturalist*, 102: 243–282.
- Santos, M. N., Oliveira, M. T., and Cúrdia, J. 2013. A comparison of the fish assemblages on natural and artificial reefs off Sal Island (Cape Verde). *Journal of the Marine Biological Association of the United Kingdom*, 93: 437–452.
- Sciberras, M., Hinz, H., Bennell, J. D., Jenkins, S. R., Hawkins, S. J., and Kaiser, M. J. 2013. Benthic community response to a scallop dredging closure within a dynamic seabed habitat. *Marine Ecology Progress Series*, 480: 83–98.
- Sheppard, C., Al-Husiani, M., Al-Jamali, F., Al-Yamani, F., Baldwin, R., Bishop, J., Benzoni, F., Dutrieux, E., Dulvy, N. K., Durvasula, S. R. V., Jones, D. A., Loughland, R., Medio, D., Nithyanandan, M., Pilling, G. M., Polikarpov, I., Price, A. R. G., Purkis, S., Riegl, B., Saburova, M., Samimi-Namin, K., Taylor, O., Wilson, S., and Zainal, K. 2010. The Gulf: a young sea in decline. *Marine Pollution Bulletin*, 60: 13–38.
- Sheppard, C., Price, A., and Roberts, C. 1992. *Marine ecology of the Arabian region: patterns and processes in extreme tropical environments*. Academic Press, London.
- Smyth, D., Al-Maslamani, I., Chatting, M., and Giraldez, B. 2016. Benthic surveys of the historic pearl oyster beds of Qatar reveal a dramatic ecological change. *Marine Pollution Bulletin*, 113: 147–155.
- Snelgrove, P., and Butman, C. 1995. Animal-sediment relationships revisited: cause versus effect. *Oceanographic Literature Review*, 8: 668.
- SPSS. 2013. *IBM SPSS Statistics for Windows*. 20. IBM Corp, Armonk, NY.
- UN. 1997. *Qatar country profile*. United Nations Commission on Sustainable Development, New York.
- UNEP. 2009. *Second National Human Development Report Qatar: Advancing Sustainable Development. Qatar National Vision 2030*. General Secretariat for Development Planning, Doha, Qatar. pp. 1–136.
- Walton, M. E. M., Al-Maslamani, I., Haddaway, N., Kennedy, H., Castillo, A., Al-Ansari, E. S., Al-Shaikh, I., Abdel-Moati, M., Al-Yafei, M. A. A., and Le Vay, L. 2016. Extreme 15N depletion in seagrasses. *Estuaries and Coasts*, 39: 1709.
- Walton, M. E. M., Al-Maslamani, I., Skov, M. W., Al-Shaikh, I., Al-Ansari, I. S., Kennedy, H. A., and Le Vay, L. 2014. Outwelling from arid mangrove systems is sustained by inwelling of seagrass productivity. *Marine Ecology Progress Series*, 507: 125–137.
- Wilson, S., Fatemi, S. M. R., Shokri, M. R., and Claereboudt, M. 2002. Status of coral reefs of the Persian/Arabian Gulf and Arabian Sea region. *In* *Status of Coral Reefs of the World*. pp. 53–62. Ed. by C. Wilkinson. Global Coral Reef Monitoring Network and Reef and Rainforest Research Centre, Townsville, Australia.
- Wu, R. S. S. 1982. Periodic defaunation and recovery in a subtropical epibenthic community, in relation to organic pollution. *Journal of Experimental Marine Biology and Ecology*, 64: 253–269.
- Yesson, C., Simon, P., Chemshirova, I., Gorham, T., Turner, C. J., Hammeken Arboe, N., Blicher, M. E., and Kemp, K. M. 2015. Community composition of epibenthic megafauna on the West Greenland Shelf. *Polar Biology*, 38: 2085–2096.

Handling editor: Joanna Norkko